

LIFTING ENDOMORPHISMS TO AUTOMORPHISMS

WILLIAM ARVESON AND DENNIS COURTNEY

ABSTRACT. Normal endomorphisms of von Neumann algebras need not be extendable to automorphisms of a larger von Neumann algebra, but they always have asymptotic lifts. We describe the structure of endomorphisms and their asymptotic lifts in some detail, and apply those results to complete the identification of asymptotic lifts of unital completely positive linear maps on von Neumann algebras in terms of their minimal dilations to endomorphisms.

1. INTRODUCTION

We work in the category whose objects are pairs (M, α) consisting of a normal unit-preserving $*$ -endomorphism $\alpha : M \rightarrow M$ of a von Neumann algebra M , and whose maps are equivariant normal $*$ -homomorphisms that map unit to unit. The isomorphisms of this category are conjugacies, in which $\alpha_1 : M_1 \rightarrow M_1$ is said to be *conjugate* to $\alpha_2 : M_2 \rightarrow M_2$ if there is a $*$ -isomorphism $\theta : M_1 \rightarrow M_2$ satisfying $\theta \circ \alpha_1 = \alpha_2 \circ \theta$.

Consider the problem of extending an endomorphism $\alpha : M \rightarrow M$ to a $*$ -automorphism of a larger von Neumann algebra, assuming that the necessary condition $\ker \alpha = \{0\}$ is satisfied. In that case α is an isometric $*$ -endomorphism of M , and a straightforward construction produces a unital C^* -algebra $N \supseteq M$ and a $*$ -automorphism β of N that restricts to α on M . This extension of α to an automorphism of a larger C^* -algebra is unique up to natural isomorphism provided one assumes that it is *minimal* in the sense that $M \cup \beta^{-1}(M) \cup \beta^{-2}(M) \cup \dots$ is norm-dense in N .

This procedure is effective for extending endomorphisms of C^* -algebras. But it is poorly suited to this category since there is no *natural* way¹ of completing the C^* -algebra N to a von Neumann algebra so as to obtain a W^* -dynamical system that extends α except in special circumstances - the most natural circumstance being that in which α preserves a faithful normal state of M . More serious problems arise when $\ker \alpha \neq \{0\}$, since in that case even extensions to C^* -algebraic automorphisms cannot exist. The proper way to associate a W^* -dynamical system to an endomorphism involves the notion of *lifting*, a concept introduced in [Arv06] for the broader category of unital completely positive maps, and which will be described momentarily.

1991 *Mathematics Subject Classification.* 46L55, 46L09.

¹It is always *possible* to carry out such a completion, but that construction does not give rise to a functor from injective endomorphisms to W^* -dynamical systems. [AK92] addresses the existence issue for E_0 -semigroups acting on von Neumann algebras.

While liftings are typically much “smaller” than the extensions of isometric endomorphisms described above, they always exist within the category and they enjoy good functorial properties.

We were led to these issues by a problem involving the broader category of normal unit-preserving completely positive linear maps $\phi : M \rightarrow M$ of von Neumann algebras M (UCP maps). It was shown in [Arv06] that every UCP map $\phi : M \rightarrow M$ has an asymptotic lift, which is unique up to natural isomorphism. Naturally, one wants to identify the asymptotic lift of ϕ in concrete terms. In [Arv06], the asymptotic lift of ϕ was identified as the tail flow of the minimal dilation of ϕ in “most” cases - namely those cases in which the tail flow of the dilated endomorphism has trivial kernel. But in general, the minimal dilation of ϕ to an endomorphism can have a nontrivial kernel, and the identification problem was left open in those cases.

The purpose of this note is to identify asymptotic lifts of UCP maps on von Neumann algebras in general. That is accomplished by first giving a description of lifts of endomorphisms, in the course of which we obtain a basic result on the structure of surjective endomorphisms of von Neumann algebras that appears to have been overlooked (Theorem 3.1). We apply these results to identify the asymptotic lift of an arbitrary UCP map in terms of its minimal dilation to an endomorphism of a larger von Neumann algebra (Theorem 4.1), thereby completing Theorem 7.1 of [Arv06].

In related work [AS06], the notion of asymptotic lift was generalized to normal positive linear maps acting on von Neumann algebras (also see [Stø06]). It is significant that since there is no dilation theory for positive linear maps that are not completely positive, the identification problem becomes a significant issue in such cases and has been only partially solved. Further discussion can be found in [AS06].

2. LIFTING ENDOMORPHISMS

Throughout this section, $\alpha : M \rightarrow M$ will denote an endomorphism acting on a von Neumann algebra M . By a W^* -dynamical system we mean a pair (N, β) , where β is a $*$ -automorphism of a von Neumann algebra N .

Definition 2.1. A *lifting* of $\alpha : M \rightarrow M$ is a triple (N, β, E) where (N, β) is a W^* -dynamical system and $E : N \rightarrow M$ is a unit-preserving normal $*$ -homomorphism satisfying $E \circ \beta = \alpha \circ E$.

Note first that for every lifting (N, β, E) of α , we have

$$(2.1) \quad E(N) \subseteq M \cap \alpha(M) \cap \alpha^2(M) \cap \cdots .$$

Indeed, every element $y = E(x)$ in the range of E can be written in the form $y = \alpha^n(E(\beta^{-n}(x))) \in \alpha^n(M)$ for every $n = 0, 1, 2, \dots$, from which the assertion is evident.

Remark 2.2 (Nondegeneracy). A lifting (N, β, E) of α is said to be *nondegenerate* if for every $x \in N$,

$$E(\beta^n(x)) = 0, \quad n \in \mathbb{Z} \implies x = 0.$$

In general, the set

$$K = \{x \in N : E(\beta^n(x)) = 0, \quad \forall n \in \mathbb{Z}\}$$

is a weak*-closed two-sided ideal in N satisfying $\beta(K) = K$. Hence there is a β -fixed central projection $c \in N$ such that $K = cN$. It follows that N decomposes into a sum $N = K \oplus N_0$, where $(N_0, \beta|_{N_0}, E|_{N_0})$ is a nondegenerate lift of α and $E(\beta^n(K)) = \{0\}$ for every $n \in \mathbb{Z}$. In particular, *every lift (N, β, E) of α can be reduced to a nondegenerate lift without affecting the range of the homomorphism E .*

Let $\alpha : M \rightarrow M$ be an endomorphism of a von Neumann algebra. The sequence of von Neumann algebras $M, \alpha(M), \alpha^2(M), \dots$ decreases as n increases, and their intersection

$$M_\infty = \bigcap_{n=1}^{\infty} \alpha^n(M)$$

is called the *tail algebra* of α . The restriction of α to the tail algebra is a *surjective* endomorphism; it is an automorphism iff $\ker \alpha \cap M_\infty = \{0\}$.

Proposition 2.3. *Let $\alpha : M \rightarrow M$ be an endomorphism. For every lifting (N, β, E) of α , the following are equivalent.*

(i) *For every normal linear functional $\rho \in M_*$, one has*

$$(2.2) \quad \lim_{n \rightarrow \infty} \|\rho \circ \alpha^n\| = \|\rho \circ E\|.$$

(ii) $E(N) = M_\infty$.

Proof. Since E is a *-homomorphism of von Neumann algebras, it maps the unit ball of N onto the unit ball of its range. Hence (ii) is equivalent to

$$E(\text{ball } N) = \text{ball } M_\infty = \bigcap_{n=1}^{\infty} \alpha^n(\text{ball } M).$$

The equivalence (i) $\iff$ (ii) now follows from the more general assertion of Lemma 3.6 of [Arv06]. $\square$

Definition 2.4. An *asymptotic lift* of an endomorphism $\alpha : M \rightarrow M$ is a nondegenerate lifting (N, β, E) satisfying the conditions of Proposition 2.3.

Remark 2.5 (Relation to asymptotic lifts of UCP maps). In [Arv06], the term *asymptotic lift* refers to a related concept that was introduced for the broader category of UCP maps on dual operator systems. It is significant that an asymptotic lift in the sense of Definition 2.4 is also an asymptotic lift in the broader sense of Definition 3.1 of [Arv06].

To prove that assertion, it suffices to show that if a lifting (N, β, E) of an endomorphism $\alpha : M \rightarrow M$ satisfies the equivalent properties (i) and

(ii) of Proposition 2.3, then those properties persist throughout the matrix hierarchy over M . Indeed, for each $n = 1, 2, \dots$, the lifting (N, β, E) of $\alpha : M \rightarrow M$ induces a natural lift $(M_n \otimes N, \text{id}_n \otimes \beta, \text{id}_n \otimes E)$ of the endomorphism $\text{id}_n \otimes \alpha : M_n \otimes M \rightarrow M_n \otimes M$, and by examining matrix entries one finds that property (ii) persists at level n . Hence property (i) holds as well for every $n = 1, 2, \dots$, and (N, β, E) satisfies Definition 3.1 of [Arv06].

Two liftings (N_k, β_k, E_k) , $k = 1, 2$, of an endomorphism $\alpha : M \rightarrow M$ are said to be *isomorphic* if there is an isomorphism of von Neumann algebras $\gamma : N_1 \rightarrow N_2$ satisfying $\gamma \circ \beta_1 = \beta_2 \circ \gamma$ and $E_2 \circ \gamma = E_1$.

Theorem 2.6. *Every endomorphism $\alpha : M \rightarrow M$ of a von Neumann algebra has an asymptotic lifting that is unique up to isomorphism.*

Proof. The proof of Theorem 3.2 of [Arv06] explicitly constructs an asymptotic lift of a UCP map of that category in terms of the space of inverse sequences of that map. Since in the present context the map is an endomorphism $\alpha : M \rightarrow M$, one sees by inspection that the constructed asymptotic lift (N, β, E) has the following properties: the space N of inverse sequences is closed under multiplication, β is a $*$ -automorphism of that von Neumann algebra, and $E : N \rightarrow M$ is a normal $*$ -homomorphism. Hence (N, β, E) is an asymptotic lift in the sense of Definition 2.4. The proof of uniqueness involves similar observations. $\square$

3. STRUCTURE OF SURJECTIVE ENDOMORPHISMS

In this section we prove that in general, a surjective endomorphism of a von Neumann algebra admits a natural decomposition into the direct sum of a W^* -dynamical system and an endomorphism of a particularly simple kind, called a *backward shift*, that depends only on $\ker \alpha$. That allows us to identify asymptotic lifts of endomorphisms in very concrete terms.

Let K be a von Neumann algebra and consider the von Neumann algebra $\ell^\infty(\mathbb{N}, K)$ of all singly-infinite bounded sequences

$$x = (x_1, x_2, \dots), \quad x_k \in K.$$

Define an endomorphism σ_+ of $\ell^\infty(\mathbb{N}, K)$ as follows

$$(3.1) \quad \sigma_+(x_1, x_2, \dots) = (x_2, x_3, \dots), \quad (x_1, x_2, \dots) \in \ell^\infty(\mathbb{N}, K).$$

Obviously, σ_+ is a normal surjective unit-preserving endomorphism, and

$$\ker \sigma_+ = K \oplus 0 \oplus 0 \oplus \dots \cong K.$$

Such an endomorphism σ_+ is called the *backward shift* based on K .

We can modify the backward shift σ_+ based on K in a nontrivial way by choosing an automorphism β of another von Neumann algebra P and letting $\beta \oplus \sigma_+$ be the endomorphism of $P \oplus \ell^\infty(\mathbb{N}, K)$ defined by

$$(3.2) \quad \beta \oplus \sigma_+ : x \oplus y \in P \oplus \ell^\infty(\mathbb{N}, K) \mapsto \beta(x) \oplus \sigma_+(y).$$

This is a surjective endomorphism whose kernel is isomorphic to K , but it has a summand P on which it restricts to an automorphism.

Theorem 3.1. *Every normal surjective endomorphism α of a von Neumann algebra is conjugate to one of the form (3.2), where σ_+ is the backward shift based on $K = \ker \alpha$, and where β is the $*$ -automorphism defined by restricting α to $P = (\mathbf{1} - c)M$, c being the α -fixed central projection*

$$(3.3) \quad c = \lim_{n \rightarrow \infty} c_n,$$

where $c_1 \leq c_2 \leq \dots$ is the sequence of central projections $\ker \alpha^n = c_n M$.

Proof. Since $\ker \alpha^n$ is a weak*-closed ideal in M for every $n = 1, 2, \dots$, it has the form $c_n M$ where c_n is a central projection; and since $\ker \alpha^n \subseteq \ker \alpha^{n+1}$, it follows that $c_n \leq c_{n+1}$.

We claim: $\alpha(c_{n+1}) = c_n$, for every $n = 1, 2, \dots$. Since $\ker \alpha^k = c_k M$ and $\alpha(M) = M$, this is equivalent to the assertion $\alpha(\ker \alpha^{n+1}) = \ker \alpha^n$. Obviously, $x \in \ker \alpha^{n+1} \implies \alpha(x) \in \ker \alpha^n$. For the opposite inclusion, choose $y \in \ker \alpha^n$. Since $M = \alpha(M)$, we can find $x \in M$ such that $y = \alpha(x)$. This x must satisfy $\alpha^{n+1}(x) = \alpha^n(\alpha(x)) = \alpha^n(y) = 0$, hence $x \in \ker \alpha^{n+1}$, and therefore $y = \alpha(x) \in \alpha(\ker \alpha^{n+1})$. These formulas $\alpha(c_{n+1}) = c_n$ clearly imply that the limit c of (3.3) is an α -fixed central projection.

Let $P = (\mathbf{1} - c)M$. Note that $\alpha(P) = \alpha(\mathbf{1} - c)\alpha(M) = (\mathbf{1} - c)M = P$. Since the kernel of α is $c_1 M \subseteq cM$ we have $P \cap \ker \alpha = \{0\}$, hence α restricts to a $*$ -automorphism of P .

Turning now to the summand cM , we claim that for every $n \geq 2$, α^{n-1} restricts to an isomorphism of von Neumann algebras

$$\alpha^{n-1} : (c_n - c_{n-1})M \cong \ker \alpha = c_1 M.$$

Indeed, the restriction of α^{n-1} to $(c_n - c_{n-1})M$ is injective because $\ker \alpha^{n-1} = c_{n-1}M$ intersects trivially with the algebra on the left. It is surjective because after iterating the formulas $\alpha(c_{k+1}) = c_k$ we find that

$$\begin{aligned} \alpha^{n-1}((c_n - c_{n-1})M) &= \alpha^{n-1}(c_n M) = \alpha^{n-1}(c_n)M = \alpha^{n-2}(c_{n-1})M \\ &= \dots = \alpha(c_2)M = c_1 M. \end{aligned}$$

Now consider the von Neumann algebra $\ell^\infty(\mathbb{N}, \ker \alpha)$, the algebra of all uniformly bounded sequences $y = (y_1, y_2, \dots)$ with $y_k \in \ker \alpha$ for $k \geq 1$. Every element $x \in cM$ admits a unique decomposition into a bounded sequence of mutually orthogonal central slices $x = x_1 + x_2 + x_3 + \dots$, where $x_1 = c_1 x$ and $x_k = (c_k - c_{k-1})x$ for $k \geq 2$. Moreover, the preceding paragraph implies that $\alpha^{k-1}(x_k) \in \ker \alpha$ for every $k \geq 2$. Thus we can define a normal homomorphism of von Neumann algebras $\theta : cM \rightarrow \ell^\infty(\mathbb{N}, \ker \alpha)$ by

$$\theta(x) = (x_1, \alpha(x_2), \alpha^2(x_3), \dots), \quad x \in cM.$$

We have also seen that for each $k \geq 2$, α^{k-1} restricts to an isomorphism from $(c_k - c_{k-1})M$ to $\ker \alpha$; and since $cM = c_1 M \oplus (c_2 - c_1)M \oplus (c_3 - c_2)M \oplus \dots$, it follows that θ is an isomorphism of von Neumann algebras.

One can now directly verify that $\theta \circ \alpha = \sigma_+ \circ \theta$, where σ_+ denotes the backward shift on $\ell^\infty(\ker \alpha)$. We conclude that θ implements a conjugacy of the restriction of α to cM and the backward shift based on $\ker \alpha$. $\square$

The W^* -dynamical system $(P, \alpha|_P)$ is called the *automorphic summand* of α . It is clear from the preceding proof that *two surjective endomorphisms are conjugate iff their automorphic summands are conjugate W^* -dynamical systems and their kernels are isomorphic von Neumann algebras*.

Theorem 3.1 leads to the following description of the asymptotic lifts of arbitrary endomorphisms of von Neumann algebras.

Corollary 3.2. *Let $\alpha : M \rightarrow M$ be an endomorphism with tail algebra $M_\infty = \bigcap_{n \geq 1} \alpha^n(M)$. Let $K = \ker \alpha \cap M_\infty$, let σ_+ be the backward shift acting on $\ell^\infty(\mathbb{N}, K)$, and let (P, β) be the automorphic summand of $\alpha|_{M_\infty}$.*

By Theorem 3.1, there is an isomorphism of von Neumann algebras

$$\theta : P \oplus \ell^\infty(\mathbb{N}, K) \rightarrow M_\infty$$

that satisfies $\theta \circ (\beta \oplus \sigma_+) = \alpha \circ \theta$. Let σ be the bilateral shift acting on the von Neumann algebra $\ell^\infty(\mathbb{Z}, K)$ by way of $\sigma(x_n) = (x_{n+1})$, and define a homomorphism $E : P \oplus \ell^\infty(\mathbb{Z}, K) \rightarrow M_\infty$ by

$$E(p \oplus (x_n)) = \theta(p \oplus (x_1, x_2, \dots)), \quad p \in P, \quad (x_n) \in \ell^\infty(\mathbb{Z}, K).$$

Then $(P \oplus \ell^\infty(\mathbb{Z}, K), \beta \oplus \sigma, E)$ is the asymptotic lift of $\alpha : M \rightarrow M$.

Proof. It is obvious that $(P \oplus \ell^\infty(\mathbb{Z}, K), \beta \oplus \sigma)$ is a W^* -dynamical system and that E is a homomorphism of von Neumann algebras with range

$$E(P \oplus \ell^\infty(\mathbb{Z}, K)) = \theta(P \oplus \ell^\infty(\mathbb{N}, K)) = M_\infty.$$

Moreover,

$$\begin{aligned} E \circ (\beta \oplus \sigma)(p \oplus (x_n)) &= E(\beta(p) \oplus (x_{n+1})) = \theta(\beta(p) \oplus (x_2, x_3, \dots)) \\ &= \theta(\beta(p) \oplus \sigma_+(x_1, x_2, \dots)) = \alpha \circ \theta(p \oplus (x_1, x_2, \dots)) \\ &= \alpha \circ E(p \oplus (x_n)), \end{aligned}$$

hence $E \circ (\beta \oplus \sigma) = \alpha \circ E$. We conclude that $(P \oplus \ell^\infty(\mathbb{Z}, K), \beta \oplus \sigma, E)$ is a lifting of α that satisfies condition (ii) of Proposition 2.3; and it remains only to show that this lifting is nondegenerate. But if $p \in P$ and $(x_n) \in \ell^\infty(\mathbb{Z}, K)$ are such that $E((\beta \circ \sigma)^k(p \oplus (x_n))) = \theta(\beta^k(p) \oplus (x_{k+1}, x_{k+2}, \dots)) = 0$ for every $k \in \mathbb{Z}$, then $\beta^k(p) = 0$ and $x_{k+1} = 0$ for every $k \in \mathbb{Z}$. The desired formula $p \oplus (x_n) = 0$ follows. $\square$

4. APPLICATION TO UCP MAPS ON VON NEUMANN ALGEBRAS

Let $\phi : M \rightarrow M$ be a UCP map acting on a von Neumann algebra M . In this section we identify the asymptotic lift of ϕ in terms of its minimal dilation to an endomorphism of a larger von Neumann algebra. This solves the identification problem in general by strengthening Theorem 7.1 of [Arv06] that was restricted to the case in which the minimal dilation has trivial kernel. Indeed, the following result applies to dilations of ϕ that are not necessarily minimal (see Chapter 8 of [Arv03]).

Theorem 4.1. *Let $\alpha : N \rightarrow N$ be an endomorphism of a von Neumann algebra and let $p \in N$ be a projection that satisfies $\alpha(p) \geq p$ and $\alpha^n(p) \uparrow \mathbf{1}$ as $n \uparrow \infty$. Let $M = pNp$ and let $\phi : M \rightarrow M$ be the UCP map defined by*

$$\phi(x) = p\alpha(x)p, \quad x \in M = pNp.$$

Let $(\tilde{N}, \tilde{\alpha}, E)$ be the asymptotic lift of α described in Corollary 3.2. Then the asymptotic lift of ϕ is $(\tilde{N}, \tilde{\alpha}, \tilde{E})$ where $\tilde{E} : \tilde{N} \rightarrow M$ is the UCP map

$$(4.1) \quad \tilde{E}(x) = pE(x)p, \quad x \in \tilde{N}.$$

Proof. Obviously $\tilde{E} : \tilde{N} \rightarrow M$ is a UCP map and we claim $\phi \circ \tilde{E} = \tilde{E} \circ \tilde{\alpha}$. Indeed, we can use $p\alpha(p) = \alpha(p)p = p$ and $\alpha \circ E = E \circ \tilde{\alpha}$ to write

$$\phi(\tilde{E}(x)) = p\alpha(\tilde{E}(x))p = p\alpha(p)\alpha(E(x))\alpha(p)p = pE(\tilde{\alpha}(x))p = \tilde{E} \circ \tilde{\alpha}(x).$$

Hence $(\tilde{N}, \tilde{\alpha}, \tilde{E})$ is a lifting of ϕ . To see that it is nondegenerate, choose $x \in \tilde{N}$ such that $\tilde{E}(\tilde{\alpha}^k(x)) = 0$, $k \in \mathbb{Z}$. Then for $n \geq 1$ we can apply α^n to $\tilde{E}(\tilde{\alpha}^{-n}(x)) = 0$ and use $\alpha \circ E = E \circ \tilde{\alpha}$ to obtain

$$0 = \alpha^n(\tilde{E}(\tilde{\alpha}^{-n}(x))) = \alpha^n(p)\alpha^n(E(\tilde{\alpha}^{-n}(x)))\alpha^n(p) = \alpha^n(p)E(x)\alpha^n(p).$$

Since $\alpha^n(p) \uparrow \mathbf{1}$ as $n \uparrow \infty$, it follows that $E(x) = 0$. Replacing x with $\tilde{\alpha}^k(x)$, $k \in \mathbb{Z}$, and using nondegeneracy of $(\tilde{N}, \tilde{\alpha}, E)$, we conclude that $x = 0$.

We claim that for every $\rho \in M_*$,

$$(4.2) \quad \lim_{n \rightarrow \infty} \|\rho \circ \phi^n\| = \|\rho \circ \tilde{E}\|.$$

To prove (4.2), fix ρ and define a normal functional $\bar{\rho} \in N_*$ by $\bar{\rho}(y) = \rho(py p)$. For every $x \in \tilde{N}$ we have $\rho \circ \tilde{E}(x) = \rho(pE(x)p) = \bar{\rho} \circ E(x)$, and as in the proof of formula (7.2) of [Arv06], we obtain the following formulas for $n \geq 1$

$$(4.3) \quad \|\rho \circ \phi^n\| = \|\bar{\rho} \circ \alpha^n\|, \quad \|\rho \circ \tilde{E}\| = \|\bar{\rho} \circ E\|.$$

Since $(\tilde{N}, \tilde{\alpha}, E)$ is the asymptotic lift of $\alpha : N \rightarrow N$, $\|\bar{\rho} \circ \alpha^n\|$ converges to $\|\bar{\rho} \circ E\|$ as $n \rightarrow \infty$, and (4.2) follows. Similarly, one can promote (4.2) throughout the matrix hierarchy over M exactly as in the proof of Theorem 7.1 of [Arv06] to complete the proof of Theorem 4.1. $\square$

REFERENCES

- [AK92] W. Arveson and A. Kishimoto. A note on extensions of E_0 -semigroups. *Proc. Amer. Math. Soc.*, 116(3):769–774, 1992.
- [Arv03] W. Arveson. *Noncommutative Dynamics and E-semigroups*. Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [Arv06] W. Arveson. The asymptotic lift of a completely positive map. *Jour. Funct. Anal.* (to appear), pages 1–23, 2006. arXiv:math.OA/0606541 v5.
- [AS06] W. Arveson and E. Størmer. Asymptotic lifts of positive linear maps. *preprint*, pages 1–13, 2006. arXiv:math.OA/0611401.
- [Stø06] E. Størmer. Multiplicative properties of positive maps. *Math. Scand.* (to appear), pages 1–8, 2006. arXiv:math.OA/0607090.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720
E-mail address: arveson@math.berkeley.edu, djc@math.berkeley.edu